Nervopex

Vitamin B₁+ B₆+ B₁₂

DESCRIPTION:

Nervopex is a combination of Vitamin B_1 (Thiamine), Vitamin B_6 (Pyridoxine) & Vitamin B_{12} (Cyanocobalamin). These vitamins play essential role as co-enzyme in the metabolism of nervous system. Thus this combination normalizes the nerve cell metabolism. **Nervopex** supports the regeneration of nerve fibers and myelin sheath by activation of the metabolism and the natural repair mechanism. The combination of these 3 Vitamins shows better efficacy than that of the individual component. In addition, Vitamin B_1 , Vitamin B_2 , combination promote body's own repair mechanism and work in pain management.

COMPOSITION:

Nervopex Tablet: Each tablet contains Thiamine Mononitrate BP 100 mg, Pyridoxine Hydrochloride BP 200 mg, and Cyanocobalamin BP 200 mg.

INDICATION:

Nervopex is indicated for the treatment of B_1 , B_6 & B_{12} deficiency syndrome. It is also indicated in the treatment of Diabetic neuropathy, Sciatica, Peripheral neuralgia, Facial paralysis, Lumbago, Intercostal neuralgia, Myalgia & Spinal pain.

DOSAGE AND ADMINISTRATION:

Nervopex Tablet: 1-2 tablets three times a day or as directed by the physician.

CONTRAINDICATION:

Nervopex is contraindicated in known hypersensitivity to any of the active ingredients in this product and in patients on Levodopa therapy.

PRECAUTION

Cyanocobalamin should not be given before a diagnosis has been fully established because of the possibility of masking symptoms of sub acute degeneration of the spinal cord. Cyanocobalamin is not a suitable form of vitamin B_{12} for the treatment of optic neuropathies associated with raised plasma concentrations of Cyanocobalamin.

USE IN PREGNANCY AND LACTATION:

Pregnancy: Nervopex is safe for pregnant women. Lactation: It is not known whether Vitamin B_1 is excreted in breast milk or not, but may be taken cautiously during lactation. Vitamin B_6 has no adverse effect during lactation. There is no known adverse effect on Vitamin B_{12} therapy also.

SIDE EFFECT:

Generally well tolerated. However, a few allergic reactions may be seen.

DRUG INTERACTION:

Levodopa: Pyridoxine (Vitamin B_{θ}) reduces therapeutic effect of levodopa but it does not occur if a dopa decarboxylase inhibitor is also given. Neomycin, Chloramphenicol, Aminosalisylic acid & histamine H_2 receptor antagonist: may reduce Vitamin B_{12} absorption.

STORAGE CONDITION:

Store in a cool dry place and keep away from light. Keep out of the reach of children.

PACKAGING:

Nervopex Tablet: Each box contains 5x10's tablets in blister pack.



Manufactured by **Apex Pharma Limited** Shafipur, Kaliakair, Gazipur, Bangladesh